arXiv:math/0212155v1 [math.QA] 11 Dec 2002

MULTIPLICATION FORMULAS FOR THE ELLIPTIC GAMMA FUNCTION

GIOVANNI FELDER* AND ALEXANDER VARCHENKO**,[1]

*Departement Mathematik, ETH-Zentrum,
8092 Zürich, Switzerland
felder@math.ethz.ch

**Department of Mathematics, University of North Carolina at Chapel Hill,
Chapel Hill, NC 27599-3250, USA
anv@email.unc.edu

Abstract. The elliptic gamma function is a generalization of the Euler gamma function. Its trigonometric and rational degenerations are the Jackson q-gamma function and the Euler gamma function. We prove multiplication formulas for the elliptic gamma function, whose degenerations are the Gauss-Askey multiplication formula for the Euler and trigonometric gamma functions.

1. Introduction

Special functions defined by infinite products often have duplication formulas. Here are some examples.

$$\sin(2\pi z) = 2\sin(\pi z)\sin(\pi(z+\frac{1}{2}))\ ,$$

$$\Gamma(2z)\sqrt{\pi} = 2^{2z-1}\,\Gamma(z)\,\Gamma(z+\frac{1}{2})\ , \tag{1}$$

$$\Gamma_q(2z)\,\Gamma_{q^2}(\frac{1}{2}) = [2]_q^{2z-1}\,\Gamma_{q^2}(z)\,\Gamma_{q^2}(z+\frac{1}{2})\ , \tag{2}$$

$$\theta_0(2z,\tau) = \theta_0(z,\tau)\,\theta_0(z+\frac{1}{2},\tau)\,\theta_0(z+\frac{\tau}{2},\tau)\,\theta_0(z+\frac{1+\tau}{2},\tau)\ . \tag{3}$$

The function $\Gamma(z)=\int_0^\infty t^{z-1}e^{-t}dt$ is the Euler gamma function. It satisfies the functional equation $\Gamma(z+1) = z\,\Gamma(z)$. Formula (1) is Legendre's duplication formula.

Date: November 2002.

[1] Supported in part by NSF grant DMS-9801582.

The function $\Gamma_q(z)$ is Jackson's q-gamma function. Set $x = e^{2\pi i z}$, $q = e^{2\pi i\tau}$, and denote

$$(x;q) \;=\; \prod_{j=0}^{\infty}(1-xq^j).$$

Then

$$\Gamma_q(z) \;=\; \Gamma_{\text{trig}}(z,\tau) \;=\; (1-q)^{1-z}\,\frac{(q;q)}{(q^z;q)}\;.$$

The q-gamma function obeys the functional equation

$$\Gamma_q(z+1) \;=\; [z]_q\;\Gamma_q(z)\;,$$

where $[z]_q = \frac{1-e^{2\pi i\tau\, z}}{1-e^{2\pi i\tau}}$ is the trigonometric analog of the number z. The q-gamma function degenerates to Euler's gamma function,

$$\lim_{\tau\to 0}\,\Gamma_{\text{trig}}(z,\tau) \;=\; \Gamma(z)\;.$$

Formula (2) is Askey's duplication formula, see [A].

The function $\theta_0(z,\tau) \;=\; (x,q)(q/x,q)$ in (3) is one of Jacobi's theta functions. Formula (3) see for instance in [Ra].

In this paper we give two duplication formulas for elliptic analogs of the gamma function,

$$\begin{aligned}\Gamma(2z,\tau,\sigma) = \quad & \Gamma(z,\tau,\sigma)\Gamma(z+\frac{\tau}{2},\tau,\sigma)\Gamma(z+\frac{\sigma}{2},\tau,\sigma)\Gamma(z+\frac{\tau+\sigma}{2},\tau,\sigma)\\ & \Gamma(z+\frac{1}{2},\tau,\sigma)\Gamma(z+\frac{1+\tau}{2},\tau,\sigma)\Gamma(z+\frac{1+\sigma}{2},\tau,\sigma)\Gamma(z+\frac{1+\tau+\sigma}{2},\tau,\sigma)\;,\end{aligned}$$

$$\bar\Gamma(2z,\tau,\sigma)\bar\Gamma(\frac{1}{2},2\tau,\sigma) = \left(\frac{\theta_0(2\tau,\sigma)}{\theta_0(\tau,\sigma)}\right)^{2z-1}\bar\Gamma(z,2\tau,\sigma)\bar\Gamma(z+\frac{1}{2},2\tau,\sigma)\;,$$

see definitions below. The expression $\frac{\theta_0(z\tau,\sigma)}{\theta_0(\tau,\sigma)}$ is an elliptic analog of the number z. We have the trigonometric and rational limits of the theta function:

$$\frac{\theta_0(z\tau,\sigma)}{\theta_0(\tau,\sigma)} \;\xrightarrow{\sigma\to i\infty}\; \frac{1-e^{2\pi i\tau\, z}}{1-e^{2\pi i\tau}} \;\xrightarrow{\tau\to 0}\; z\;.$$

2. ELLIPTIC GAMMA FUNCTION

The *elliptic gamma function* is an elliptic generalization of the Euler gamma function. It is the meromorphic function of three complex variables z,τ,σ, with $\operatorname{Im}\tau, \operatorname{Im}\sigma > 0$ defined by the convergent infinite product

$$\Gamma(z,\tau,\sigma) = \prod_{j,k=0}^{\infty}\frac{1-e^{2\pi i((j+1)\tau+(k+1)\sigma-z)}}{1-e^{2\pi i(j\tau+k\sigma+z)}}\;.$$

It is the unique solution of a functional equation involving the Jacobi theta function θ_0.

Theorem. [FV1] *Suppose that* τ, σ *are complex numbers with positive imaginary part. Then* $u(z) = \Gamma(z, \tau, \sigma)$ *is the unique meromorphic solution of the difference equation*

$$u(z + \sigma) = \theta_0(z, \tau)u(z)$$

such that:

(i) $u(z)$ *obeys* $u(z + 1) = u(z)$ *and is holomorphic on the upper half plane* $\operatorname{Im} z > 0$,
(ii) $u((\tau + \sigma)/2) = 1$.

The elliptic gamma function first appeared in [R]. The modular properties of the elliptic gamma function and their relations to $SL(3, \mathbb{Z})$ are discussed in [FV1], appearances and applications of the elliptic gamma function can be found in [B, DP, JMN, JKKMW, FTV, FV2, FV3, FV4].

Let $\bar{\Gamma}$ be the function

$$\bar{\Gamma}(z, \tau, \sigma) \;=\; \frac{(q; q)}{(r; r)}\, \theta_0(\tau, \sigma)^{1-z}\, \Gamma(z\tau, \tau, \sigma)\ , \qquad q = e^{2\pi i\tau},\ r = e^{2\pi i\sigma}.$$

Then $u(z) = \bar{\Gamma}(z, \tau, \sigma)$ is a solution of the functional equation

$$u(z + 1) \;=\; \frac{\theta_0(\tau z, \sigma)}{\theta_0(\tau, \sigma)}\, u(z).$$

The normalization was chosen here so that $u(1) = 1$. As $\sigma \to i\infty$ we recover Jackson's q-gamma function,

$$\Gamma_{\text{trig}}(z, \tau) \;=\; \lim_{\sigma \to i\infty} \bar{\Gamma}(z, \tau, \sigma)\ .$$

3. Multiplication Formulas

3.1. The first multiplication formula.

Theorem. *For any natural* n *we have*

$$\Gamma(nz, \tau, \sigma) \;=\; \prod_{k_1, k_2, k_3 = 0}^{n-1} \Gamma(z + \frac{k_1 + k_2\tau + k_3\sigma}{n}, \tau, \sigma)\ .$$

Proof. Let $w = e^{2\pi i/n}$. Then the right hand side of this formula is

$$\prod_{k_1,k_2,k_3=0}^{n-1} \prod_{l,m=0}^{\infty} \frac{1 - w^{-k_1} q^{l+\frac{n-k_2}{n}} r^{m+\frac{n-k_3}{n}} x^{-1}}{1 - w^{k_1} q^{l+\frac{k_2}{n}} r^{m+\frac{k_3}{n}} x}$$

$$= \prod_{k_2,k_3=0}^{n-1} \prod_{l,m=0}^{\infty} \frac{1 - q^{(l+1)n-k_2} r^{(m+1)n-k_3} x^{-n}}{1 - q^{ln+k_2} r^{mn+k_3} x^n} = \prod_{l,m=0}^{\infty} \frac{1 - q^{l+1} r^{m+1} x^{-n}}{1 - q^l r^m x^n} = \Gamma(nz, \tau, \sigma).$$

□

3.2. The second multiplication formula.

Theorem. *For any natural n we have*

$$\bar\Gamma(nz,\tau,\sigma)\ \bar\Gamma(\frac{1}{n},n\tau,\sigma)\ \bar\Gamma(\frac{2}{n},n\tau,\sigma)\ \dots\ \bar\Gamma(\frac{n-1}{n},n\tau,\sigma)\ =$$
$$\left(\frac{\theta_0(n\tau,\sigma)}{\theta_0(\tau,\sigma)}\right)^{nz-1} \bar\Gamma(z,n\tau,\sigma)\ \bar\Gamma(z+\frac{1}{n},n\tau,\sigma)\ \bar\Gamma(z+\frac{2}{n},n\tau,\sigma)\ \dots\ \bar\Gamma(z+\frac{n-1}{n},n\tau,\sigma)\ .$$

The theorem is an easy consequence of the following two lemmas.

Lemma. *For any natural m and n, we have*

$$\Gamma(z,\tau,\sigma)\ =\ \prod_{a=0}^{m-1}\prod_{b=0}^{n-1} \Gamma(z+a\tau+b\sigma, m\tau, n\sigma)\ .$$

Lemma. *For any natural n, we have*

$$\Gamma(\tau,n\tau,\sigma)\ \Gamma(2\tau,n\tau,\sigma)\ \dots\ \Gamma((n-1)\tau,n\tau,\sigma)\ =\ \frac{1}{(q,q^n)(q^2,q^n)\dots(q^{n-1},q^n)}\ .$$

References

[A] R. Askey, *The q-gamma and q-beta functions*, Applicable Anal. 8 (1978/79), no. 2, 125–141.

[B] R. J. Baxter, *Partition function of the eight-vertex lattice model*, Ann. Phys., 70(1972) 193–228

[DP] B. Davies and I. Peschel, *A unified treatment of Ising model magnetizations*, Ann. Physik (1997) 187–214

[FTV] G. Felder, V. Tarasov and A. Varchenko, *Solutions of the elliptic qKZB equations and Bethe ansatz I*, Amer. Math. Soc. Transl. 180 (1997), 45–75; *Monodromy of solutions of the elliptic Knizhnik-Zamolodchikov-Bernard difference equations*, q-alg/9705017

[FV1] G. Felder and A. Varchenko, *The Elliptic Gamma Function and $SL(3,\mathbb{Z}) \ltimes \mathbb{Z}^3$*, Advances in Math. 156 (2000), 44–76.

[FV2] G. Felder and A. Varchenko, *The q-deformed Knizhnik-Zamolodchikov-Bernard heat equation*, Comm. Math. Phys. 221 (2001), no. 3, 549–571;

[FV3] G. Felder and A. Varchenko, *q-deformed KZB heat equation: completeness, modular properties and SL(3,Z)*, QA/0110081 .

[FV4] G. Felder and A. Varchenko, *Even powers of divisors and elliptic zeta values*, math.QA/0205116.

[JMN] M. Jimbo, T. Miwa and A. Nakayashiki, *Difference equations for the correlation functions of the eight-vertex model*, J. Phys. A: Math. Gen. 26 (1993) 2199–2209

[JKKMW] M. Jimbo, R. Kedem, H. Konno, T. Miwa and R. Weston, *Difference equations in spin chains with a boundary*, Nucl. Phys. B 448 [FS] (1995) 429–456

[Ra] E. Rainville, *Special Functions*, Macmillan Co., 1960.

[R] S. N. M. Ruijsenaars, *First order analytic difference equations and integrable quantum systems*, J. Math. Phys. 38 (1997), no. 2, 1069–1146